

SINGULAR LOCI OF GRASSMANN-HIBI TORIC VARIETIES

J. BROWN AND V. LAKSHMIBAI[†]

ABSTRACT. For the toric variety X associated to the Bruhat poset of Schubert varieties in the Grassmannian, we describe the singular locus in terms of the faces of the associated polyhedral cone. We further show that the singular locus is pure of codimension 3 in X , and the generic singularities are of cone type. We also determine the tangent cones at the maximal singularities of X . These turn out to be again toric varieties. In the case of X being associated to the Bruhat poset of Schubert varieties in the Grassmannian of 2-planes in a n -dimensional vector space (over the base field), we also prove a certain product formula relating the multiplicities at certain singular points.

INTRODUCTION

Let K denote the base field which we assume to be algebraically closed of arbitrary characteristic. Given a distributive lattice $\mathcal{L}$, let $X(\mathcal{L})$ denote the affine variety in $\mathbb{A}^{\#\mathcal{L}}$ whose vanishing ideal is generated by the binomials $X_\tau X_\varphi - X_{\tau \vee \varphi} X_{\tau \wedge \varphi}$ in the polynomial algebra $K[X_\alpha, \alpha \in \mathcal{L}]$ (here, $\tau \vee \varphi$ (resp. $\tau \wedge \varphi$) denotes the *join* - the smallest element of $\mathcal{L}$ greater than both τ, φ (resp. the *meet* - the largest element of $\mathcal{L}$ smaller than both τ, φ)). These varieties were extensively studied by Hibi in [13] where Hibi proves that $X(\mathcal{L})$ is a normal variety. On the other hand, Eisenbud-Sturmfels show in [8] that a binomial prime ideal is toric (here, “toric ideal” is in the sense of [20]). Thus one obtains that $X(\mathcal{L})$ is a normal toric variety. We shall refer to such a $X(\mathcal{L})$ as a *Hibi toric variety*.

For $\mathcal{L}$ being the Bruhat poset of Schubert varieties in a minuscule G/P , it is shown in [10] that $X(\mathcal{L})$ flatly deforms to $\widehat{G/P}$ (the cone over G/P), i.e., there exists a flat family over $\mathbb{A}^1$ with $\widehat{G/P}$ as the generic fiber and $X(\mathcal{L})$ as the special fiber. More generally, for a Schubert variety $X(w)$ in a minuscule G/P , it is shown in [10] that $X(\mathcal{L}_w)$ flatly deforms to $\widehat{X(w)}$, the cone over $X(w)$ (here, $\mathcal{L}_w$ is the Bruhat poset of Schubert subvarieties of $X(w)$). In a subsequent paper (cf. [11]), the authors of loc.cit., studied the singularities of $X(\mathcal{L})$, $\mathcal{L}$ being the Bruhat poset of Schubert varieties in the Grassmannian; further, in loc.cit., the authors gave a conjecture (cf. §11 in loc.cit; see also Remark 9.1 of this paper for a statement of this conjecture) giving a necessary and sufficient condition for a point on $X(\mathcal{L})$ to be smooth, and

[†] Partially supported by NSF grant DMS-0400679 and NSA-MDA904-03-1-0034.

proved in loc.cit., the sufficiency part of the conjecture. Subsequently, the necessary part of the conjecture was proved in [2] by Batyrev et al. The toric varieties $X(\mathcal{L})$, $\mathcal{L}$ being the Bruhat poset of Schubert varieties in the Grassmannian play an important role in the area of mirror symmetry; for more details, see [2, 3]. We refer to such an $X(\mathcal{L})$ as a *Grassmann-Hibi toric variety* (abbreviated *G-H toric variety*).

The proof (in [11]) of the sufficiency part of the conjecture of [11] uses the Jacobian criterion for smoothness, while the proof (in [2]) of the necessary part of the conjecture of [11] uses certain desingularization of $X(\mathcal{L})$.

In this paper, our first main result is a description of the singular locus of a G-H toric variety $X(\mathcal{L})$ in terms of the faces of the associated polyhedral cone; in particular, we give a proof of the conjecture of [11] using just the combinatorics of the polyhedral cone associated to the toric variety $X(\mathcal{L})$. Further, we prove (cf. Theorem 6.19) that the singular locus of $X(\mathcal{L})$ is pure of codimension 3 in $X(\mathcal{L})$, and that the generic singularities are of cone type (more precisely, the singularity type is the same as that at the vertex of the cone over the quadric surface $x_1x_4 - x_2x_3 = 0$ in $\mathbb{P}^3$); we also determine (in loc.cit) the tangent cone to $X(\mathcal{L})$ at a maximal singularity which turns out to be a toric variety (here, by a “maximal singularity”, we mean a point $P \in X(\mathcal{L})$ such that the closure of the torus orbit through P is an irreducible component of $\text{Sing } X(\mathcal{L})$, the singular locus of $X(\mathcal{L})$). We further obtain (cf. Corollary 7.3, Theorem 7.11) an interpretation of the multiplicities at some of the singularities as certain Catalan numbers in the case of $\mathcal{L}$ being the Bruhat poset of Schubert varieties in the Grassmannian of 2-planes in K^n . We also present a product formula (cf. Theorem 7.17).

It turns out that the conjecture of [11] does not extend to a general $X(\mathcal{L})$ (see §9.2 for a counter example). But, recently (cf. [6]), we have proved the conjecture of [11] for other minuscule posets. It should be remarked that this paper contains more results for the Grassmann-Hibi toric varieties which can not be deduced from the results of [6].

The singularities of the Hibi toric variety were also studied by Wagner (cf. [21]).

The sections are organized as follows: In §1, we recall generalities on toric varieties. In §2, we recall generalities on distributive lattices. In §3, we introduce the Hibi toric variety $X(\mathcal{L})$ and recall some results from [11, 17] on $X(\mathcal{L})$. In §4, we recall results from [17] on the polyhedral cone associated to $X(\mathcal{L})$. In §5, we introduce the Grassmann-Hibi toric variety. In §6, we prove our first main result giving the description of $\text{Sing } X(\mathcal{L})$ in terms of faces of the cone associated to $X(\mathcal{L})$; and we present our results on the tangent cones and deduce the multiplicities at the associated points. In §7, we present our results for $X(\mathcal{L})$, $\mathcal{L}$ being the Bruhat poset of Schubert varieties in the Grassmannian of 2-planes in K^n . In §8, we present a formula for the multiplicity at the unique T -fixed point of $X(\mathcal{L})$, $\mathcal{L}$ being the Bruhat poset of Schubert varieties in the Grassmannian of d -planes in K^n . In §9, we present a counter

example to show that the conjecture of [11] does not extend to a general $X(\mathcal{L})$; in this section, we also present two conjectures.

1. GENERALITIES ON TORIC VARIETIES

Since our main object of study is a certain affine toric variety, we recall in this section some basic definitions on affine toric varieties. Let $T = (K^*)^m$ be an m -dimensional torus.

Definition 1.1. (cf. [9], [15]) An *equivariant affine embedding* of a torus T is an affine variety $X \subseteq \mathbb{A}^l$ containing T as an open subset and equipped with a T -action $T \times X \rightarrow X$ extending the action $T \times T \rightarrow T$ given by multiplication. If in addition X is normal, then X is called an *affine toric variety*.

1.2. The Cone Associated to a Toric Variety. Let M be the character group of T , and N the $\mathbb{Z}$ -dual of M . Recall (cf. [9], [15]) that there exists a strongly convex rational polyhedral cone $\sigma \subset N_{\mathbb{R}} (= N \otimes_{\mathbb{Z}} \mathbb{R})$ such that

$$K[X] = K[S_{\sigma}],$$

where S_{σ} is the subsemigroup $\sigma^{\vee} \cap M$, $\sigma^{\vee}$ being the cone in $M_{\mathbb{R}}$ dual to σ . Note that S_{σ} is a finitely generated subsemigroup in M .

1.3. Orbit Decomposition in Affine Toric Varieties. We shall denote X also by X_{σ} . We may suppose, without loss of generality, that σ spans $N_{\mathbb{R}}$ so that the dimension of σ equals $\dim N_{\mathbb{R}} = \dim T$. (Here, by dimension of σ , one means the vector space dimension of the span of σ .)

Let us first recall the definition of faces of a convex polyhedral cone:

Definition 1.4. A *face* τ of σ is a convex polyhedral sub cone of σ of the form $\tau = \sigma \cap u^{\perp}$ for some $u \in \sigma^{\vee}$, and is denoted $\tau < \sigma$. Note that σ itself is considered a face.

We have that X_{τ} is a principal open subset of X_{σ} , namely,

$$X_{\tau} = (X_{\sigma})_u$$

Each face τ determines a (closed) point P_{τ} in X_{σ} , namely, it is the point corresponding to the maximal ideal in $K[X] (= K[S_{\sigma}])$ given by the kernel of $e_{\tau} : K[S_{\sigma}] \rightarrow K$, where for $u \in S_{\sigma}$, we have

$$e_{\tau}(u) = \begin{cases} 1, & \text{if } u \in \tau^{\perp} \\ 0, & \text{otherwise} \end{cases}$$

Remark 1.5. As a point in $\mathbb{A}^l$, P_{τ} may be identified with the l -tuple with 1 at the i -th place if χ_i is in $\tau^{\perp}$, and 0 otherwise (here, χ_i denotes the weight of the T -weight vector y_i - the class of x_i in $K[X_{\sigma}]$).

1.6. Orbit Decomposition. Let O_τ denote the T -orbit in X_σ through P_τ . We have the following orbit decomposition in X_σ :

$$\begin{aligned} X_\sigma &= \bigcup_{\theta \leq \sigma} O_\theta \\ \overline{O_\tau} &= \bigcup_{\theta \geq \tau} O_\theta \\ \dim \tau + \dim O_\tau &= \dim X_\sigma \end{aligned}$$

See [9], [15] for details.

Thus $\tau \mapsto \overline{O_\tau}$ defines an order reversing bijection between $\{\text{faces of } \sigma\}$ and $\{T\text{-orbit closures in } X_\sigma\}$. In particular, we have the following two extreme cases:

1. If τ is the 0-face, then $P_\tau = (1, \dots, 1)$ (as a point in $\mathbb{A}^l$), and $O_\tau = T$, and is contained in X_θ , for every $\theta < \sigma$. It is a dense open orbit.
2. If $\tau = \sigma$, then $P_\tau = (0, \dots, 0)$ (as a point in $\mathbb{A}^l$), and $O_\tau = \{P_\tau\}$, the unique closed orbit. (Note that since σ spans $N_\mathbb{R}$, P_σ is a T -fixed point and is in fact the unique T -fixed point in X_σ .)

For a face τ , let us denote by N_τ the sublattice of N generated by the lattice points of τ . Let $N(\tau) = N/N_\tau$, and $M(\tau)$, the $\mathbb{Z}$ -dual of $N(\tau)$. For a face θ of σ such that θ contains τ as a face, set

$$\theta_\tau := (\theta + (N_\tau)_\mathbb{R}) / (N_\tau)_\mathbb{R}.$$

Then the collection $\{\theta_\tau, \sigma \geq \theta \geq \tau\}$ gives the set of faces of the cone $\sigma_\tau (\subset N(\tau)_\mathbb{R})$.

Lemma 1.7 (cf. [9]). *For a face $\tau < \sigma$, $\overline{O_\tau}$ gets identified with the toric variety $\text{Spec } K[S_{\sigma_\tau}]$. Further, $K[\overline{O_\tau}] = K[S_\sigma \cap \tau^\perp]$.*

2. TORIC VARIETIES ASSOCIATED TO FINITE DISTRIBUTIVE LATTICES

We shall now study a special class of toric varieties, namely, the toric varieties associated to distributive lattices. We shall first collect some definitions on finite partially ordered sets. A partially ordered set is also called a poset.

Definition 2.1. A finite poset P is called *bounded* if it has a unique maximal, and a unique minimal element, denoted $\hat{1}$ and $\hat{0}$ respectively.

Definition 2.2. A totally ordered subset C of a finite poset P is called a *chain*, and the number $\#C - 1$ is called the *length* of the chain.

Definition 2.3. A bounded poset P is said to be *graded* (or also *ranked*) if all maximal chains have the same length (note that $\hat{1}$ and $\hat{0}$ belong to any maximal chain).

Definition 2.4. Let P be a graded poset. The length of a maximal chain in P is called the *rank* of P .

Definition 2.5. Let P be a graded poset. For $\lambda, \mu \in P$ with $\lambda \geq \mu$, the graded poset $\{\tau \in P \mid \mu \leq \tau \leq \lambda\}$ is called the *interval from μ to λ* , and denoted by $[\mu, \lambda]$. The rank of $[\mu, \lambda]$ is denoted by $l_\mu(\lambda)$; if $\mu = \hat{0}$, then we denote $l_\mu(\lambda)$ by just $l(\lambda)$.

Definition 2.6. Let P be a graded poset, and $\lambda, \mu \in P$, with $\lambda \geq \mu$. The ordered pair (λ, μ) is called a *cover* (and we also say that λ *covers* μ) if $l_\mu(\lambda) = 1$.

2.7. Generalities on distributive lattices.

Definition 2.8. A *lattice* is a partially ordered set $(\mathcal{L}, \leq)$ such that, for every pair of elements $x, y \in \mathcal{L}$, there exist elements $x \vee y$ and $x \wedge y$, called the *join*, respectively the *meet* of x and y , defined by:

$$\begin{aligned} x \vee y &\geq x, \quad x \vee y \geq y, \quad \text{and if } z \geq x \text{ and } z \geq y, \text{ then } z \geq x \vee y, \\ x \wedge y &\leq x, \quad x \wedge y \leq y, \quad \text{and if } z \leq x \text{ and } z \leq y, \text{ then } z \leq x \wedge y. \end{aligned}$$

It is easy to check that the operations $\vee$ and $\wedge$ are commutative and associative.

Definition 2.9. Given a lattice $\mathcal{L}$, a subset $\mathcal{L}' \subset \mathcal{L}$ is called a *sublattice* of $\mathcal{L}$ if $x, y \in \mathcal{L}'$ implies $x \wedge y \in \mathcal{L}'$, $x \vee y \in \mathcal{L}'$.

Definition 2.10. Two lattices $\mathcal{L}_1$ and $\mathcal{L}_2$ are *isomorphic* if there exists a bijection $\varphi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ such that, for all $x, y \in \mathcal{L}_1$,

$$\varphi(x \vee y) = \varphi(x) \vee \varphi(y) \quad \text{and} \quad \varphi(x \wedge y) = \varphi(x) \wedge \varphi(y).$$

Definition 2.11. A lattice is called *distributive* if the following identities hold:

- (1) $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
- (2) $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z).$

Definition 2.12. An element z of a lattice $\mathcal{L}$ is called *join-irreducible* (respectively *meet-irreducible*) if $z = x \vee y$ (respectively $z = x \wedge y$) implies $z = x$ or $z = y$. The set of join-irreducible (respectively meet-irreducible) elements of $\mathcal{L}$ is denoted by $J(\mathcal{L})$ (respectively $M(\mathcal{L})$), or just by J (respectively M) if no confusion is possible.

Definition 2.13. The set $J(\mathcal{L}) \cap M(\mathcal{L})$ of join and meet-irreducible elements is denoted by $JM(\mathcal{L})$, or just JM if no confusion is possible.

Definition 2.14. A subset I of a poset P is called an *ideal* of P if for all $x, y \in P$,

$$x \in I \text{ and } y \leq x \text{ imply } y \in I.$$

Theorem 2.15 (Birkhoff). *Let $\mathcal{L}$ be a distributive lattice with $\hat{0}$, and P the poset of its nonzero join-irreducible elements. Then $\mathcal{L}$ is isomorphic to the lattice of ideals of P , by means of the lattice isomorphism*

$$\alpha \mapsto I_\alpha = \{\tau \in P \mid \tau \leq \alpha\}, \quad \alpha \in \mathcal{L}.$$

Definition 2.16. A quadruple of the form $(\tau, \phi, \tau \vee \phi, \tau \wedge \phi)$, with $\tau, \phi \in \mathcal{L}$ non-comparable is called a *diamond*, and is denoted by $D(\tau, \phi, \tau \vee \phi, \tau \wedge \phi)$ or also just $D(\tau, \phi)$. The unordered pair (τ, ϕ) (respectively $(\tau \vee \phi, \tau \wedge \phi)$) is called the *skew* (respectively *main*) diagonal of the diamond $D(\tau, \phi)$.

The following Lemma is easily checked.

Lemma 2.17. *With the notations as above, we have*

- (a) $J = \{\tau \in \mathcal{L} \mid \text{there exists at most one cover of the form } (\tau, \lambda)\}.$
- (b) $M = \{\tau \in \mathcal{L} \mid \text{there exists at most one cover of the form } (\lambda, \tau)\}.$

For $\alpha \in \mathcal{L}$, let I_α be the ideal corresponding to α under the isomorphism in Theorem 2.15; (we will also use the notation $I(\alpha)$).

Lemma 2.18 (cf. [17]). *Let (τ, λ) be a cover in $\mathcal{L}$. Then I_τ equals $I_\lambda \dot{\cup} \{\beta\}$ for some $\beta \in J(\mathcal{L})$.*

3. THE VARIETY $X(\mathcal{L})$

Consider the polynomial algebra $K[X_\alpha, \alpha \in \mathcal{L}]$; let $I(\mathcal{L})$ be the ideal generated by $\{X_\alpha X_\beta - X_{\alpha \vee \beta} X_{\alpha \wedge \beta}, \alpha, \beta \in \mathcal{L}\}$. Then one knows (cf.[13]) that $K[X_\alpha, \alpha \in \mathcal{L}] / I(\mathcal{L})$ is a normal domain; in particular, we have that $I(\mathcal{L})$ is a prime ideal. Let $X(\mathcal{L})$ be the affine variety of the zeroes in K^l of $I(\mathcal{L})$ (here, $l = \#\mathcal{L}$). Then $X(\mathcal{L})$ is an affine normal variety defined by binomials; on the other hand, by [8], a binomial prime ideal is a toric ideal (here, “toric ideal” is in the sense of [20]). Hence $X(\mathcal{L})$ is a toric variety for the action by a suitable torus T .

In the sequel, we shall denote $R(\mathcal{L}) = K[X_\alpha, \alpha \in \mathcal{L}] / I(\mathcal{L})$. Further, for $\alpha \in \mathcal{L}$, we shall denote the image of X_α in $R(\mathcal{L})$ by x_α .

Definition 3.1. The variety $X(\mathcal{L})$ will be called a *Hibi toric variety*.

Remark 3.2. An extensive study of $X(\mathcal{L})$ appears first in [13].

We have that $\dim X(\mathcal{L}) = \dim T$.

Theorem 3.3 (cf. [17]). *The dimension of $X(\mathcal{L})$ is equal to $\#J(\mathcal{L})$.*

Definition 3.4. For a finite distributive lattice $\mathcal{L}$, we call the cardinality of $J(\mathcal{L})$ the *dimension* of $\mathcal{L}$, and we denote it by $\dim \mathcal{L}$. If $\mathcal{L}'$ is a sublattice of $\mathcal{L}$, then the *codimension* of $\mathcal{L}'$ in $\mathcal{L}$ is defined as $\dim \mathcal{L} - \dim \mathcal{L}'$.

Definition 3.5. (cf. [21]) A sublattice $\mathcal{L}'$ of $\mathcal{L}$ is called an *embedded sublattice* of $\mathcal{L}$ if

$$\tau, \phi \in \mathcal{L}, \quad \tau \vee \phi, \tau \wedge \phi \in \mathcal{L}' \quad \Rightarrow \quad \tau, \phi \in \mathcal{L}'.$$

Given a sublattice $\mathcal{L}'$ of $\mathcal{L}$, let us consider the variety $X(\mathcal{L}')$, and consider the canonical embedding $X(\mathcal{L}') \hookrightarrow \mathbb{A}(\mathcal{L}') \hookrightarrow \mathbb{A}(\mathcal{L})$ (here $\mathbb{A}(\mathcal{L}') = \mathbb{A}^{\#\mathcal{L}'}$, $\mathbb{A}(\mathcal{L}) = \mathbb{A}^{\#\mathcal{L}}$).

Proposition 3.6 (cf. [11]). *$X(\mathcal{L}')$ is a subvariety of $X(\mathcal{L})$ if and only if $\mathcal{L}'$ is an embedded sublattice of $\mathcal{L}$.*

3.7. Degeneration of $X(\mathcal{L})$ to a monomial scheme.

Definition 3.8. A monomial $x_{\alpha_1} \cdots x_{\alpha_r}$ in $R(\mathcal{L})$ of degree r is said to be *standard* if $\alpha_1 \geq \cdots \geq \alpha_r$.

Theorem 3.9 (cf. [11]). *Monomials standard on $X(\mathcal{L})$ form a basis for $R(\mathcal{L})$.*

Definition 3.10. Let $S(\mathcal{L})$ denote the *Stanley-Reisner algebra* of $\mathcal{L}$, namely, $K[X_{\alpha}, \alpha_i \in \mathcal{L}] / \mathfrak{b}(\mathcal{L})$ where $\mathfrak{b}(\mathcal{L})$ is the (monomial ideal) ideal in $K[X_{\alpha}, \alpha_i \in \mathcal{L}]$ generated by $\{X_{\alpha}X_{\beta}, \alpha, \beta \text{ incomparable}\}$.

Note that $\text{Spec } S(\mathcal{L})$ is a monomial scheme. Using the standard monomial basis, a flat deformation of $\text{Spec } S(\mathcal{L})$ is constructed in [11]:

Theorem 3.11. *There exists a flat family over $\mathbb{A}^1$ whose special fiber ($t = 0$) is $\text{Spec } S(\mathcal{L})$, and generic fiber (t invertible) is $X(\mathcal{L})$.*

3.12. Hilbert polynomial and Hilbert series: Recall the Hilbert function of a graded, finitely generated K -algebra $R = \bigoplus_m R_m$; $\phi_m(R) := \dim R_m$ is the *Hilbert function* of R . Recall that for $m \gg 0$, $\phi_m(R)$ is given by $P(m)$, where $P(x)$ is a polynomial of degree equal to $r := \dim R - 1 (= \dim \text{Proj } R)$ and coefficients are in $\mathbb{Q}$; further, the leading coefficient of $P(x)$ is of the form $\frac{e_R}{r!}$. The polynomial $P(x)$ is called *the Hilbert polynomial* of R , and e_R is called the *degree of R* or also the *degree of $\text{Proj } R$* . The series $H_R(t) := \sum \phi_m t^m$ is called *the Hilbert series of R* .

Tangent cone and multiplicity: Let $X = \text{Spec } B \hookrightarrow \mathbb{A}^l$ be an affine variety; let S be the polynomial algebra $K[X_1, \dots, X_l]$. Let $P \in X$, and let M_P be the maximal ideal in $K[X]$ corresponding to P (we are concerned only with closed points of X). Let $A = \mathcal{O}_{X,P}$, the stalk at P ; denote the unique maximal ideal in A by $M (= M_P B_{M_P})$. Then $\text{Spec } gr(A, M)$, where $gr(A, M) = \bigoplus_{j \in \mathbb{Z}_+} M^j / M^{j+1}$ is the *tangent cone* to X at P , and is denoted $TC_P X$. Note that

$$gr(B, M_P) = gr(A, M)$$

The degree of the graded ring $gr(A, M)$ is defined as the *multiplicity of X at P* , and is denoted $\text{mult}_P X$

Fact: With notation as above, if B is graded, then taking P to be $\mathbf{0}$ (the origin), we have a natural identification

$$TC_{\mathbf{0}} X \cong X$$

Hence in this case, we have

$$\text{mult}_{\mathbf{0}} X = e_B$$

See [7] for details.

Square-free monomial schemes: Let $A = K[X_1, \dots, X_n]$. Let I be a square-free monomial ideal in A ; i.e., generated by a bunch of square-free monomials $\mathbf{m}_i, i \in J$,

for some finite indexing set J ; let $R = A/I$. Let

$$I_i = \{j \mid X_j \text{ occurs in } \mathbf{m}_i\}.$$

Theorem 3.13 (cf. [17]).

- (1) $\phi_m(R) = \sum \binom{m-1}{\#S-1}$, the sum running over all S such that $S \not\supseteq I_i, \forall i$. In particular, the Hilbert polynomial of R has degree $d-1$, where $d = \max \{\#S, S \not\supseteq I_i, \forall i\}$.
- (2) $H_R(t) = \sum \frac{t^{\#S}}{(1-t)^{\#S}}$, the sum running over all S such that $S \not\supseteq I_i, \forall i$.
- (3) $\dim \operatorname{Spec} R = d$.
- (4) $\deg R = \#\{S, S \not\supseteq I_i, \forall i, \#S = d\}$

Corollary 3.14. *Let P be a ranked poset, and $R(P)$ be the Stanley-Reisner algebra of P (cf. Definition 3.10). Then $\dim \operatorname{Spec} R(P)$ equals the cardinality of the set of elements in a maximal chain in P , and $\deg R(P)$ is equal to the number of maximal chains in P .*

Now returning to our situation of $X(\mathcal{L})$, we get, in view of Theorem 3.11, and the above Corollary

Theorem 3.15.

- (1) *The dimension of $X(\mathcal{L})$ equals the cardinality of the set of elements in a maximal chain in $\mathcal{L}$*
- (2) *The degree of $K[X(\mathcal{L})]$ is equal to the number of maximal chains in $\mathcal{L}$.*

Combining the above Theorem with Theorem 3.3, and Lemma 2.18, we obtain

Theorem 3.16.

- (1) *The cardinality of $J(\mathcal{L})$ equals the cardinality of the set of elements in a maximal chain in $\mathcal{L}$.*
- (2) *Fix any maximal chain $\beta_1 < \dots < \beta_d$ in $X(\mathcal{L})$, d being $\#J(\mathcal{L})$. Let γ_{i+1} be the element of $J(\mathcal{L})$ corresponding to the cover (β_{i+1}, β_i) , $i \geq 1$ (cf. Lemma 2.18); set $\gamma_1 = \beta_1$. Then $J(\mathcal{L}) = \{\gamma_1, \dots, \gamma_d\}$*

Now $X(\mathcal{L})$ being of cone type, i.e, the vanishing ideal of $X(\mathcal{L})$ (as a subvariety of $\mathbb{A}^{\# \mathcal{L}}$) is homogeneous, we have (in view of §3.12, Fact) the following

Theorem 3.17.

- (1) *We have an identification of $X(\mathcal{L})$ with $TC_{P_\sigma} X(\mathcal{L})$ (the tangent cone to $X(\mathcal{L})$ at P_σ , origin being identified with P_σ)*
- (2) *$\operatorname{mult}_{P_\sigma} X(\mathcal{L})$ equals the degree of $K[X(\mathcal{L})]$ ($= \#\{\text{maximal chains in } \mathcal{L}\}$).*

4. CONE AND DUAL CONE OF $X(\mathcal{L})$:

Denote the poset of join-irreducibles by $J(\mathcal{L})$. Denote by $\mathcal{I}$ the poset of ideals of $J(\mathcal{L})$. For $A \in \mathcal{I}$, denote by $\mathbf{m}_A$ the monomial:

$$\mathbf{m}_A := \prod_{\tau \in A} y_\tau$$

in the polynomial algebra $K[y_\tau, \tau \in J(\mathcal{L})]$. If α is the element of $\mathcal{L}$ such that $I_\alpha = A$ (cf. Theorem 2.15), then we shall denote $\mathbf{m}_A$ also by $\mathbf{m}_\alpha$. Consider the surjective algebra map

$$F : K[X_\alpha, \alpha \in \mathcal{L}] \rightarrow K[\mathbf{m}_A, A \in \mathcal{I}], \quad X_\alpha \mapsto \mathbf{m}_A, \quad A = I_\alpha$$

Theorem 4.1 (cf. [13], [17]). *We have an isomorphism $K[X(\mathcal{L})] \cong K[\mathbf{m}_A, A \in \mathcal{I}]$.*

Let us denote the torus acting on the toric variety $X(\mathcal{L})$ by T ; note that $\dim T = \#J(\mathcal{L}) = d$, say. Identifying T with $(K^*)^d$, let $\{f_z, z \in J(\mathcal{L})\}$ denote the standard $\mathbb{Z}$ -basis for $X(T)$, namely, for $t = (t_z, z \in J(\mathcal{L}))$, $f_z(t) = t_z$. Denote $M := X(T)$; let N be the $\mathbb{Z}$ -dual of M , and $\{e_y, y \in J(\mathcal{L})\}$ be the basis of N dual to $\{f_z, z \in J(\mathcal{L})\}$. For $A \in \mathcal{I}$, set

$$f_A := \sum_{z \in A} f_z$$

Let $V = N_{\mathbb{R}} (= N \otimes_{\mathbb{Z}} \mathbb{R})$. Let $\sigma \subset V$ be the cone such that $X(\mathcal{L}) = X_\sigma$. Let $\sigma^\vee \subset V^*$ be the cone dual to σ . Let $S_\sigma = \sigma^\vee \cap M$, so that $K[X(\mathcal{L})]$ equals the semi group algebra $K[S_\sigma]$.

As an immediate consequence of Theorem 4.1, we have

Proposition 4.2. *The semigroup S_σ is generated by $f_A, A \in \mathcal{I}$.*

Let $M(J(\mathcal{L}))$ be the set of maximal elements in the poset $J(\mathcal{L})$. Let $Z(J(\mathcal{L}))$ denote the set of all covers in the poset $J(\mathcal{L})$ (i.e., $(z, z'), z > z'$ in the poset $J(\mathcal{L})$, and there is no other element $y \in J(\mathcal{L})$ such that $z > y > z'$). For a cover $(y, y') \in Z(J(\mathcal{L}))$, denote

$$v_{y, y'} := e_{y'} - e_y$$

Proposition 4.3 (cf. [17]). *The cone σ is generated by $\{e_z, z \in M(J(\mathcal{L}))\}$, $v_{y, y'}, (y, y') \in Z(J(\mathcal{L}))\}$.*

4.4. Analysis of faces of σ . We shall concern ourselves just with the closed points in $X(\mathcal{L})$. So in the sequel, by a point in $X(\mathcal{L})$, we shall mean a closed point. Let τ be a face of σ . Let P_τ be the distinguished point of O_τ with the associated maximal ideal being the kernel of the map

$$K[S_\sigma] \rightarrow K, \\ u \in S_\sigma, u \mapsto \begin{cases} 1, & \text{if } u \in \tau^\perp \\ 0, & \text{otherwise} \end{cases}$$

Then for a point $P \in X(\mathcal{L})$ (identified with a point in $\mathbb{A}^l$, $l = \#\mathcal{L}$), denoting by $P(\alpha)$, the α -th co-ordinate of P , we have,

$$P_\tau(\alpha) = \begin{cases} 1, & \text{if } f_{I_\alpha} \in \tau^\perp \\ 0, & \text{otherwise} \end{cases}$$

With notation as above, let

$$D_\tau = \{\alpha \in \mathcal{L} \mid P_\tau(\alpha) \neq 0\}$$

We have,

4.5. The bijection $\mathcal{D}$. (cf. [17]) We have a bijection

$$\mathcal{D} : \{\text{faces of } \sigma\} \xleftrightarrow{bij} \{\text{embedded sublattices of } \mathcal{L}\}, \quad \mathcal{D}(\tau) = D_\tau$$

Proposition 4.6. [cf. [17]] *Let τ be a face of σ . Then we have $\overline{O_\tau} = X(D_\tau)$.*

5. THE DISTRIBUTIVE LATTICE $I_{d,n}$ AND THE GRASSMANN-HIBI TORIC VARIETY

We now turn our focus to

$$\mathcal{L} = I_{d,n} = \{x = (i_1, \dots, i_d) \mid 1 \leq i_1 < \dots < i_d \leq n\}.$$

The partial order $\geq$ on $I_{d,n}$ is given by

$$(i_1, \dots, i_d) \geq (j_1, \dots, j_d) \iff i_1 \geq j_1, \dots, i_d \geq j_d.$$

For $x \in I_{d,n}$, we denote the j -th entry in x by $x(j)$, $1 \leq j \leq d$.

Lemma 5.1. $I_{d,n}$ is a distributive lattice.

Proof. To see that $I_{d,n}$ is a lattice, we simply need to see that there is a well defined meet and join. We see that

$$(i_1, \dots, i_d) \vee (j_1, \dots, j_d) = (M_1, \dots, M_d),$$

where $M_k = \max\{i_k, j_k\}$. It's clear that this is an element of the lattice. Similarly,

$$(i_1, \dots, i_d) \wedge (j_1, \dots, j_d) = (m_1, \dots, m_d),$$

for $m_k = \min\{i_k, j_k\}$. For proof of distributivity, see [14]. □

Remark 5.2. It is a well known fact (see [16] for example) that the partially ordered set $I_{d,n}$ is isomorphic to the poset determined by the set of Schubert varieties in $G_{d,n}$, the Grassmannian of d -dimensional subspaces in an n -dimensional space, the Schubert varieties being partially ordered by inclusion.

In the lemma below, by a *segment* we shall mean a set consisting of consecutive integers.

Lemma 5.3 (cf. [11]). *We have*

- (a) *The element $\tau = (i_1, \dots, i_d)$ is join-irreducible if and only if either τ is a segment, or τ consists of two disjoint segments (μ, ν) , with μ starting with 1.*
- (b) *The element $\tau = (i_1, \dots, i_d)$ is meet-irreducible if and only if either τ is a segment, or τ consists of two disjoint segments (μ, ν) , with ν ending with n .*
- (c) *The element $\tau = (i_1, \dots, i_d)$ is join-irreducible and meet-irreducible if and only if either τ is a segment, or τ consists of two disjoint segments (μ, ν) , with μ starting with 1 and ν ending with n .*

Definition 5.4. We shall denote $X(I_{d,n})$ by just $X_{d,n}$ and will refer to it as a *Grassmann-Hibi toric variety*, or a *G-H toric variety* for short.

6. SINGULAR FACES OF THE G-H TORIC VARIETY $X_{d,n}$

Let $\mathcal{L}$ represent the distributive lattice $I_{d,n}$. Let $J(\mathcal{L})$ be the join irreducible elements of $\mathcal{L}$. From Lemma 5.3, we have that the elements of $J(\mathcal{L})$ are of two types, elements of Type I consist of segments of $\mathcal{L}$, and those of Type II consist of all elements $\tau \in \mathcal{L}$ such that $\tau = (\mu, \nu)$ where μ is a segment beginning at 1 and ν is just a segment.


Since the generators of the cone σ are determined by $J(\mathcal{L})$ (cf. Proposition 4.3), we will often consider $J(\mathcal{L})$ as a partially ordered set with the partial order induced from $\mathcal{L}$. Notice that $J(\mathcal{L})$ has one maximal element, which is also the maximal element of $\mathcal{L}$: $\hat{1} = (n - d + 1, \dots, n)$; and $J(\mathcal{L})$ has one minimal element, which is also the minimal element of $\mathcal{L}$: $\hat{0} = (1, \dots, d)$. Viewing $J(\mathcal{L})$ as a poset, for each element x of $J(\mathcal{L})$, there are at most two covers of the form (y, x) .

For example, if $x = (1, \dots, k, l + 1, \dots, l + d - k) \in J(\mathcal{L})$, we have $y = (1, \dots, k, l + 2, \dots, l + d - k + 1)$ and $y' = (1, \dots, k - 1, l, \dots, l + d - k)$ forming the two covers of x in $J(\mathcal{L})$ (if $k = 1$, then $y' = (l, \dots, l + d - 1)$). If $l = n - d + k$, or if x is of Type I, then x has only one cover.

Lemma 6.1 (cf. [18]). *The partially ordered set $J(\mathcal{L})$ is a distributive lattice.*

Remark 6.2. As a lattice, $J(\mathcal{L})$ looks like a tessellation of diamonds in the shape of a rectangle, with sides of length $d - 1$ and $n - d - 1$. For example, let $d = 3$ and

$n = 7$. Then $J(\mathcal{L})$ is the lattice



The Face σ_{ij} : As in §4, let σ be the cone associated to $X(\mathcal{L})$.

Definition 6.3. For $1 \leq i \leq n - d - 1$, $1 \leq j \leq d - 1$, let

$$\mu_{ij} = (1, \dots, j, i + j + 1, \dots, i + d), \text{ and}$$

$$\lambda_{ij} = (i + 1, \dots, i + j, n + 1 + j - d, \dots, n).$$

Define

$$\mathcal{L}_{ij} = \mathcal{L} \setminus [\mu_{ij}, \lambda_{ij}].$$

Remark 6.4. (i) By [11], Lemma 11.5, we have that $\mathcal{L}_{ij}$ is an embedded sublattice.
(ii) For $\alpha, \beta \in J(\mathcal{L})$, $\alpha \wedge \beta = \mu_{ij}$ for some $1 \leq i \leq n - d - 1$, $1 \leq j \leq d - 1$; thus every diamond in $J(\mathcal{L})$ has a μ_{ij} as its minimal element.

Definition 6.5. Let σ_{ij} be the face of σ for which $D_{\sigma_{ij}} = \mathcal{L}_{ij}$.

Definition 6.6. A face τ of σ is a *singular* (resp. *non-singular*) face if P_τ is a singular (resp. non-singular) point of X_σ .

Our first result is that σ_{ij} is a singular face. To prove this, we first determine a set of generators for σ_{ij} .

Definition 6.7. Let us denote by $W(\sigma)$ (or just W) the set of generators for σ as described in Proposition 4.3. For a face τ of σ , define

$$W(\tau) = \{v \in W \mid f_{I(\alpha)}(v) = 0, \forall \alpha \in D_\tau\}.$$

(Here, D_τ is as in §4.5). Then $W(\tau)$ gives a set of generators for τ .

6.8. Determination of $W(\sigma_{ij})$. It will aid our proof below to notice a few facts about the generators of σ_{ij} . First of all, $e_{\hat{1}}$ is not a generator for any σ_{ij} ; because $\hat{1} \in \mathcal{L}_{ij}$ for all $1 \leq i \leq n-d-1$, $1 \leq j \leq d-1$, and $e_{\hat{1}}$ is non-zero on $f_{I_{\hat{1}}}$. Similarly, for the cover $(y', \hat{0})$, where $y' = (1, \dots, d-1, d+1)$; $e_{\hat{0}} - e_{y'}$ is not a generator for any σ_{ij} .

Secondly, for any cover (y', y) in $J(\mathcal{L})$, $e_y - e_{y'}$ is not a generator of σ_{ij} if $y \in \mathcal{L}_{ij}$, because $f_{I(y)}(e_y - e_{y'}) \neq 0$. Thus, in determining elements of $W(\sigma_{ij})$, we need only be concerned with elements $e_y - e_{y'}$ such that $y \in J(\mathcal{L}) \cap [\mu_{ij}, \lambda_{ij}]$. The elements of $J(\mathcal{L}) \cap [\mu_{ij}, \lambda_{ij}]$ are

$$\begin{aligned} y_t &= (1, \dots, j, i+j+1+t, \dots, i+d+t) \text{ for } 0 \leq t \leq n-d-i \\ z_t &= (1, \dots, j-t, i+j+1-t, \dots, i+d) \text{ for } 0 \leq t \leq j \end{aligned}$$

Note that $y_0 = z_0 = \mu_{ij}$, $z_j = (i+1, \dots, i+d)$. We shall now prove in the Theorem below that $W(\sigma_{ij})$ consists of precisely four elements, forming a diamond in the distributive lattice $J(\mathcal{L})$ with μ_{ij} as the smallest element.

Theorem 6.9. $W(\sigma_{ij}) = \{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$, where A , B , and C are defined in the proof.

Proof. We divide the proof into two cases: $j = 1$, and $j > 1$.

Case 1: Let $j = 1$ and $1 \leq i \leq n-d-1$. Here we have

$$\mu_{ij} = (1, i+2, \dots, i+d) \text{ and } \lambda_{ij} = (i+1, n-d+2, \dots, n).$$

As discussed previously, we find that μ_{ij} is covered in $J(\mathcal{L})$ by $A = (i+1, \dots, i+d)$ and $B = (1, i+3, \dots, i+d+1)$. We have that both A and B are in the interval $[\mu_{ij}, \lambda_{ij}]$. Let C be the join of A and B in the lattice $J(\mathcal{L})$, i.e.

$$C = (i+2, \dots, i+d+1).$$

Note that (C, A) and (C, B) are covers in $J(\mathcal{L})$.

We first observe the following:

(*) If $x = (x_1, \dots, x_d) \in \mathcal{L}_{ij}$ is $\geq \mu_{ij}$, then $x \geq C$

(this follows since the facts that $x \geq \mu_{ij}$ and $x \in \mathcal{L}_{ij}$ imply that $x \not\leq \lambda_{ij}$, and hence $x_1 \geq i+2$).

Claim (i): $e_{\mu_{ij}} - e_A$ and $e_{\mu_{ij}} - e_B$ are both in $W(\sigma_{ij})$.

We shall prove the claim for $e_{\mu_{ij}} - e_A$ (the proof for $e_{\mu_{ij}} - e_B$ is similar). To prove that $e_{\mu_{ij}} - e_A$ is in $W(\sigma_{ij})$, we need to show that there does not exist $x = (x_1, \dots, x_d) \in \mathcal{L}_{ij}$ such that $x \geq \mu_{ij}$ and $x \not\geq A$. But this follows from (*) (note that (*) implies that if $x = (x_1, \dots, x_d) \in \mathcal{L}_{ij}$ is $\geq \mu_{ij}$, then $x \geq A$).

Claim (ii): $e_A - e_C$ and $e_B - e_C$ are in $W(\sigma_{ij})$.

The proof is similar to that of Claim (i). Again, we shall prove the result for $e_A - e_C$, (the proof for $e_B - e_C$ is similar). We need to show that there does not exist $x = (x_1, \dots, x_d) \in \mathcal{L}_{ij}$ such that $x \geq A$, but $x \not\geq C$. Again this follows from (*) (note that $x \geq A$ implies in particular that $x \geq \mu_{ij}$).

Claim (iii): $W(\sigma_{ij}) = \{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$.

In the case under consideration, j being 1, the only elements of $J(\mathcal{L}) \cap [\mu_{ij}, \lambda_{ij}]$ are of the form

$$y_t = (1, i+t+2, \dots, i+d+t) \text{ for } 0 \leq t \leq n-d-i;$$

$$\text{and } z_1 = (i+1, \dots, i+d).$$

Let $y'_t = (i+t+1, \dots, i+d+t)$ for $1 \leq t \leq n-d-i$; thus we have covers of the type (y'_t, y_t) for $0 \leq t \leq n-d-i$, and (y_{t+1}, y_t) for $0 \leq t \leq n-d-1-i$. Note that $y_0 = \mu_{ij}$, $y_1 = B$, $z_1 = y'_0 = A$ and $y'_1 = C$. In Claims (i) and (ii), we have shown that the covers (y_1, y_0) , (y'_0, y_0) , (y'_1, y_1) , and (y'_1, z_1) yield elements of $W(\sigma_{ij})$. Note also that C is the only cover of A . Hence, it only remains to show that $e_{y_t} - e_{y'_t} \notin W(\sigma_{ij})$ for $2 \leq t \leq n-d-i$, and $e_{y_t} - e_{y_{t+1}} \notin W(\sigma_{ij})$ for $1 \leq t \leq n-d-1-i$. For each of these covers, we shall exhibit a $x \in \mathcal{L}_{ij}$ such that $f_{I(x)}$ is non-zero on the cover under consideration.

Define $x_t = (i+t, i+t+2, \dots, i+d+t)$; we have that $x_t \in \mathcal{L}_{ij}$ for $2 \leq t \leq n-d-i$. Further, $f_{I(x_t)}$ is non-zero on $e_{y_t} - e_{y_{t+1}}$ for $2 \leq t \leq n-d-i-1$ and $e_{y_t} - e_{y'_t}$ for $2 \leq t \leq n-d-i$. For (y_2, y_1) , note that $C \in \mathcal{L}_{ij}$, and $f_{I(C)}$ is non-zero on $e_{y_1} - e_{y_2}$. This completes the proof of Case 1.

Case 2: Now let $2 \leq j \leq d-1$, $1 \leq i \leq n-d-1$. We have

$$\mu_{ij} = (1, \dots, j, i+j+1, \dots, i+d), \text{ and}$$

$$\lambda_{ij} = (i+1, \dots, i+j, n+1+j-d, \dots, n).$$

As in Case 1, we will look for covers of μ_{ij} in $J(\mathcal{L})$. They are $A = (1, \dots, j-1, i+j, \dots, i+d)$, and $B = (1, \dots, j, i+j+2, \dots, i+d+1)$. Define C to be the join of A and B in the lattice $J(\mathcal{L})$, thus

$$C = (1, \dots, j-1, i+j+1, \dots, i+d+1).$$

Claim (iv): $\{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$ are in $W(\sigma_{ij})$.

We first observe the following:

(**) If $x = (x_1, \dots, x_d) \in \mathcal{L}_{ij}$ is $\geq \mu_{ij}$, then $x \geq C$

For, suppose that $x \not\geq C$. Now the facts that $x \geq \mu_{ij}$ and $x \in \mathcal{L}_{ij}$ imply that $x \not\geq \lambda_{ij}$, and thus $x_l > i+l$ for some $1 \leq l \leq j$. Also, $x \not\geq C$; hence $x_k < i+k+1$ for some $j \leq k \leq d$. Therefore

$$x = (x_1, \dots, x_{l-1}, x_l > i+l, x_{l+1} > i+l+1, \dots, x_{k-1} > i+k-1,$$

$$i+k+1 > x_k > i+k, \dots)$$

Clearly, no such x_k exists, and thus (**) follows.

By (**), we have that if $x \in \mathcal{L}_{ij}$ is such that $x \geq \mu_{ij}$, then $x \geq A, B$, and C . Hence Claim (iv) follows.

Claim (v): $W(\sigma_{ij}) = \{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$.

As in Claim (iii), we will show that all other covers in $J(\mathcal{L})$ of the form (y', y) where $y \in J(\mathcal{L}) \cap [\mu_{ij}, \lambda_{ij}]$ are not in $W(\sigma_{ij})$. As in §6.8, all of the elements of $J(\mathcal{L}) \cap [\mu_{ij}, \lambda_{ij}]$ are

$$\begin{aligned} y_t &= (1, \dots, j, i+j+1+t, \dots, i+d+t) \text{ for } 0 \leq t \leq n-d-i \\ z_t &= (1, \dots, j-t, i+j+1-t, \dots, i+d) \text{ for } 0 \leq t \leq j \end{aligned}$$

(note that $z_j = (i+1, \dots, i+d)$). We will examine covers of these elements; note that $y_0 = z_0 = \mu_{ij}$, $z_1 = A$, and $y_1 = B$.

Let $z'_t = (1, \dots, j-t, i+j+2-t, \dots, i+d+1)$ for $1 \leq t \leq n-d-i$, and $z'_j = (i+2, \dots, i+d+1)$. First, we want to show that the covers $(z_{t+1}, z_t)_{1 \leq t \leq j-1}$ and $(z'_t, z_t)_{2 \leq t \leq j}$ do not yield elements in $W(\sigma_{ij})$. Note that $(z'_1, z_1) = (C, A)$, and $e_A - e_C \in W(\sigma_{ij})$. Also, $C \in \mathcal{L}_{ij}$, and $f_{I(C)}(e_{z_1} - e_{z_2})$ is non-zero; thus $e_{z_1} - e_{z_2} \notin W(\sigma_{ij})$, and we may restrict our concern to $t \geq 2$. Let

$$\begin{aligned} x_t &= (1, \dots, j-t, i+j+1-t, n-d+j-t+2, \dots, n) \text{ for } 2 \leq t \leq j-1, \\ x_j &= (i+1, n-d+2, \dots, n). \end{aligned}$$

Now, on the interval $2 \leq t \leq j-1$, we have the following facts:

- (1) $x_t \geq z_t$,
- (2) $x_t \not\geq z_{t+1}$,
- (3) $x_t \not\geq z'_t$.
- (4) $x_t \not\geq \lambda_{ij}$

Facts (1), (3), and (4) above hold for the case $t = j$, it is just a separate check. Therefore, for $2 \leq t \leq j$ (resp. $2 \leq t \leq j-1$), we have $x_t \in \mathcal{L}_{ij}$ and $f_{I(x_t)}$ is non-zero on $e_{z_t} - e_{z'_t}$ (resp. $e_{z_t} - e_{z_{t+1}}$).

Next, we must concern ourselves with covers involving y_t . Define

$$y'_t = (1, \dots, j-1, i+j+t, \dots, i+d+t) \text{ for } 1 \leq t \leq n-d-i.$$

To complete Claim (v), we must show that the covers

$$(y_{t+1}, y_t)_{1 \leq t \leq n-d-i-1} \text{ and } (y'_t, y_t)_{2 \leq t \leq n-d-i}$$

do not yield elements of $W(\sigma_{ij})$. Note that $(y'_1, y_1) = (C, B)$ and thus does yield an element of $W(\sigma_{ij})$. Also, $f_{I(C)}(e_{y_1} - e_{y_2})$ is non-zero, therefore we can restrict our attention to $t \geq 2$. Let $x'_t = (1, \dots, j-1, i+j+1, i+j+t+1, \dots, i+d+t)$. On the interval $2 \leq t \leq n-d-i$, we have the following facts:


- (1) $x'_t \geq y_t$,
- (2) $x'_t \not\geq y'_t$,

- (3) $x'_t \not\geq y_{t+1}$, (for $t \leq n - d - i - 1$),
- (4) $x'_t \not\leq \lambda_{ij}$.

Therefore, on the interval $2 \leq t \leq n - d - i$ (resp. $2 \leq t \leq n - d - i - 1$), we have $x'_t \in \mathcal{L}_{ij}$ and $f_{I(x'_t)}$ is non-zero on $e_{y_t} - e_{y'_t}$ (resp. $e_{y_t} - e_{y_{t+1}}$).

This completes Claim (v), Case 2, and the proof of the theorem. $\square$

Remark 6.10. The face σ_{ij} corresponds to the diamond in $J(\mathcal{L})$:



which is a poset of rank 2.

Lemma 6.11. $\dim(\sigma_{ij}) = 3$.

Proof. We have a set of generators for σ_{ij} , namely $\{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$. We can see that a subset of three of these generators is linearly independent. Thus if the fourth generator can be put in terms of the first three, the result follows. Notice that

$$(e_{\mu_{ij}} - e_A) - (e_{\mu_{ij}} - e_B) + (e_A - e_C) = e_B - e_C.$$

$\square$

As an immediate consequence of Theorem 6.9 and Lemma 6.11, we have the following

Theorem 6.12. *We have an identification of the (open) affine piece in $X(\mathcal{L})$ corresponding to the face $\sigma_{i,j}$ with the product $Z \times (K^*)^{\#J(\mathcal{L})-3}$, where Z is the cone over the quadric surface $x_1x_4 - x_2x_3 = 0$ in $\mathbb{P}^3$.*

We now prove two lemmas which hold for a general toric variety.

Lemma 6.13. *Let X_τ be an affine toric variety with τ as the associated cone. Then X_τ is a non-singular variety if and only if it is non-singular at the distinguished point P_τ .*

Proof. Only the implication $\Leftarrow$ requires a proof. Let then P_τ be a smooth point. Let us assume (if possible) that $\text{Sing } X_\tau \neq \emptyset$. We have the following facts:

- $\text{Sing } X_\tau$ is a closed T -stable subset of X_τ .
- $P_\tau \in \overline{O_\theta}$, for every face θ of τ (see §1.6); in particular, $P_\tau \in \overline{O_\theta}$, for some face θ such that P_θ is a singular point, (such a θ exists, since by our assumption $\text{Sing } X_\tau$ is non-empty). Therefore we obtain that $P_\tau \in \text{Sing } X_\tau$, a contradiction. Hence our assumption is wrong and the result follows. $\square$

Lemma 6.14. *Let τ be a face of σ . Then P_τ is a smooth point of X_σ if and only if P_τ is a smooth point of X_τ , i.e., if and only if τ is generated by a part of a basis of N (N is the $\mathbb{Z}$ dual of the character group of the torus).*

Proof. We have that X_τ is a principal open subset of X_σ . Hence X_σ is non-singular at P_τ if and only if X_τ is non-singular at P_τ . By Lemma 6.13, X_τ is non-singular at P_τ if and only if X_τ is a non-singular variety; but by §2.1 of [9], this is true if and only if τ is generated by a part of a basis of N . $\square$

Theorem 6.15. *Let $\tau = \sigma_{i,j}$. We have*

- (1) $P_\tau \in \text{Sing } X_\sigma$.
- (2) *We have an identification of $TC_{P_\tau} X_\sigma$ with $Z \times (K)^{\#J(\mathcal{L})-3}$, Z being as in Theorem 6.12; in particular, $TC_{P_\tau} X_\sigma$ is a toric variety.*
- (3) *The singularity at P_τ is of the same type as that at the vertex of the cone over the quadric surface $x_1x_4 - x_2x_3 = 0$ in $\mathbb{P}^3$. In particular, $\text{mult}_{P_\tau} X_\sigma = 2$.*

Proof. Assertion (1) follows from Lemmas 6.13, 6.14 and Theorem 6.12.

X_τ being open in X_σ , we may identify $TC_{P_\tau} X_\sigma$ with $TC_{P_\tau} X_\tau$; assertion (2) follows from this in view of Theorem 6.12 (and §3.12).

Assertion (3) is immediate from (2). $\square$

Next, we will show that the faces containing some σ_{ij} are the only singular faces. We first prove some preparatory Lemmas.

Lemma 6.16. *Let $A \neq \hat{0}$. If $e_A - e_C$ is in W (the set of generators of σ as described in Proposition 4.3), then it is in $W(\sigma_{ij})$ (cf. Definition 6.7) for some (i, j) where $1 \leq i \leq n - d - 1$, $1 \leq j \leq d - 1$.*

Proof. If A is equal to some μ_{ij} , then C must be one of the two covers of $\mu_{ij}(= (1, \dots, j, i + j + 1, \dots, i + d))$ in $J(\mathcal{L})$, and we are done by Theorem 6.9. Thus, we will assume that $A \neq \mu_{ij}$. Hence A is a join irreducible of one of the following two forms.

Case 1: $A = (1, \dots, k, n - d + k + 1, \dots, n)$ for some k .

Then $\mu_{n-d-1,k} = (1, \dots, k, n - d + k, \dots, n - 1)$; and $(A, \mu_{n-d-1,k})$ is a cover in $J(\mathcal{L})$. Also, A only has one cover in $J(\mathcal{L})$, which must be C , thus $e_A - e_C$ is an element of $W(\sigma_{n-d-1,k})$ as shown in Cases 1 and 2 of Theorem 6.9.

Case 2: $A = (k + 1, \dots, k + d)$, $1 \leq k \leq n - d - 1$ (note that $k < n - d$, since $C > A$ because $e_A - e_C \in W$).

Then we have $\mu_{k,1} = (1, k + 2, \dots, k + d)$; and $(A, \mu_{k,1})$ is a cover in $J(\mathcal{L})$. Also, we must have $C = (k + 2, \dots, k + d + 1)$, and $e_A - e_C$ is an element of $W(\sigma_{k,1})$ by Case 1 of Theorem 6.9. $\square$

We now return to the case of a G-H toric variety.

Theorem 6.17. *Let τ be a face such that D_τ is not contained in any $\mathcal{L}_{ij}$, $1 \leq i \leq n-d-1$, $1 \leq j \leq d-1$. Then the associated face τ is nonsingular (i.e., if any face τ does not contain any one σ_{ij} , then τ is nonsingular).*

Proof. By Lemma 6.14, for τ to be nonsingular, it must be generated by part of a basis for N . Since τ is generated by a subset $W(\tau)$ of W , for τ to be singular its generators would have to be linearly dependent. (Generally this is not enough to prove a face is singular or nonsingular, but since all generators in W have coefficients equal to ± 1 , any linearly independent set will serve as part of a basis for N .) Suppose τ is singular; then there is some subset of the elements of $W(\tau)$ equal to $\{e_1 - e_2, \dots\}$ such that $\sum a_{ij}(e_i - e_j) = 0$, with coefficients a_{ij} nonzero for at least one (i, j) .

Recall that the elements of W can be represented as all the line segments in the lattice $J(\mathcal{L})$, with the exception of e_1 (see diagram in Remark 6.2). Therefore, the linearly dependent generators of τ must represent a “loop” of line segments in $J(\mathcal{L})$. This loop will have at least one bottom corner, left corner, top corner, and right corner.

Choose some particular $\mathcal{L}_{ij}$. From the previous section, we have that $W(\sigma_{ij}) = \{e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B, e_A - e_C, e_B - e_C\}$. These four generators are represented by the four sides of a diamond in $J(\mathcal{L})$. Thus, by hypothesis, the generators of τ represent a loop in $J(\mathcal{L})$ that does not traverse all four sides of the diamond representing all four generators of σ_{ij} .

Since by hypothesis, D_τ is not contained in any $\mathcal{L}_{ij}$, $1 \leq i \leq n-d-1$, $1 \leq j \leq d-1$, there must be at least one element of D_τ in the interval $[\mu_{ij}, \lambda_{ij}]$, say $\alpha \in [\mu_{ij}, \lambda_{ij}]$. We have, $\alpha \geq \mu_{ij}$ and $\alpha \not\geq C$ for C as defined in the proof of Theorem 6.9. Based on how α compares to both A and B , we can eliminate certain elements of W from $W(\tau)$. There are four possibilities; we list all four, as well as the corresponding generators in $W(\sigma_{ij})$ which are not in $W(\tau)$, i.e., those generators v in $W(\sigma_{ij})$ such that $f_{I(\alpha)}(v) \neq 0$:

$$\begin{aligned} \alpha \not\geq A, \alpha \not\geq B &\Rightarrow e_{\mu_{ij}} - e_A, e_{\mu_{ij}} - e_B \notin W(\tau) \\ \alpha \geq A, \alpha \not\geq B &\Rightarrow e_A - e_C, e_{\mu_{ij}} - e_B \notin W(\tau) \\ \alpha \not\geq A, \alpha \geq B &\Rightarrow e_{\mu_{ij}} - e_A, e_B - e_C \notin W(\tau) \\ \alpha \geq A, \alpha \geq B &\Rightarrow e_A - e_C, e_B - e_C \notin W(\tau) \end{aligned}$$

Therefore, it is impossible to have $\{e_{\mu_{ij}} - e_A, e_A - e_C\}$ or $\{e_{\mu_{ij}} - e_B, e_B - e_C\}$ contained in $W(\tau)$. This is true for any (i, j) ; therefore (in view of Lemma 6.16) our “loop” in $J(\mathcal{L})$ that represented the generators of τ cannot have a left corner or a right corner. Therefore it is really not possible to have a loop at all, and so the generators of τ are linearly independent, and the result follows. $\square$

Corollary 6.18. *The G - H toric variety $X_{d,n}$ is smooth along the orbit O_τ if and only if the face τ does not contain any σ_{ij} .*

Combining the above Theorem with Theorem 6.15 and Lemma 6.11, we obtain our first main Theorem:

Theorem 6.19. *Let $\mathcal{L} = I_{d,n}$. Then*


- (1) $\text{Sing } X(\mathcal{L}) = \bigcup_{\sigma_{i,j}} \overline{O}_{\sigma_{i,j}}$, the union being taken over all $\sigma_{i,j}$'s as in Theorem 6.9.
- (2) $\text{Sing } X(\mathcal{L})$ is pure of codimension 3 in $X(\mathcal{L})$; further, the generic singularities are of cone type (more precisely, the singularity type is the same as that at the vertex of the cone over the quadric surface $x_1x_4 - x_2x_3 = 0$ in $\mathbb{P}^3$).
- (3) For $\tau = \sigma_{i,j}$, $TC_{P_\tau} X(\mathcal{L})$ is a toric variety; further, $\text{mult}_{P_\tau} X(\mathcal{L}) = 2$.

Remark 6.20. Thus Theorem 6.19 proves the conjecture of [11] using just the combinatorics of the cone associated to the toric variety $X_{d,n}$ (for a statement of the conjecture of [11], see Remark 9.1). Further, it gives a description of $\text{Sing } X_{d,n}$ purely in terms of the faces of the cone associated to $X_{d,n}$.

7. MULTIPLICITIES OF SINGULAR FACES OF $X_{2,n}$

In this section, we take $\mathcal{L} = I_{2,n}$, determine the multiplicity of $X_{2,n}(= X(I_{2,n}))$ at P_τ for certain of the singular faces of $X_{2,n}$, and deduce a product formula. For $I_{d,n}$, we defined $\mathcal{L}_{ij}$ and the corresponding face $\sigma_{i,j}$ for $1 \leq j \leq d-1$, $1 \leq i \leq n-d-1$; thus for $I_{2,n}$, we need to consider only $\mathcal{L}_{i,1}$ for $1 \leq i \leq n-3$.

For example, below is the poset of join irreducibles for $I_{2,6}$. We write $\sigma_{i,1}$ inside each diamond because the four segments surrounding it represent the four generators of the face.



To go from the join irreducibles of $I_{2,6}$ to $I_{2,7}$, we just add $(1, 7)$ and $(6, 7)$ to the poset above, forming $\sigma_{4,1}$. We will see that this makes the calculation of the multiplicities of singular faces of $I_{2,n}$ much easier.

In the sequel, we shall denote the set of join irreducibles of $I_{2,n}$ by $J_{2,n}$; also, as in the previous sections, σ will denote the cone corresponding to $X_{2,n}$.

7.1. $\text{Mult}_{P_\sigma} X_{2,n}$. Let σ be the CPC associated to $I_{d,n}$. Now $X_{d,n}$ being of cone type (i.e., the vanishing ideal is homogeneous), we have a canonical identification of $T_{P_\sigma} X_{d,n}$ (the tangent cone to $X_{d,n}$ at P_σ) with $X_{d,n}$. Hence by Theorem 3.17, (2), we have that $\text{mult}_{P_\sigma} X_{d,n}$ equals the number of maximal chains in $I_{d,n}$. So we begin by counting the number of maximal chains in $I_{2,n}$.

As we move through a chain from $(1, 2)$, at any point (i, j) we have at most two possibilities for the next point: $(i + 1, j)$ or $(i, j + 1)$. For each cover in our chain, we assign a value: for a cover of type $((i, j + 1), (i, j))$ assign $+1$; for a cover of type $((i + 1, j), (i, j))$ assign -1 .

A maximal chain C in $I_{2,n}$ contains $2n - 3$ lattice points, and thus every chain can be uniquely represented by a $(2n - 4)$ -tuple of 1 's and -1 's; let us denote this $(2n - 4)$ -tuple by $n_C = \langle a_1, \dots, a_{2n-4} \rangle$.

What rules do we have on n_C ? First it is clear that 1 and -1 occur precisely $n - 2$ times. Secondly, we can see that $a_1 = +1$, and for any $1 \leq k \leq 2n - 4$, if $\{a_1, \dots, a_k\}$ contains more -1 's than $+1$'s, then we have arrived at a point (i, j) with $i > j$, which is not a lattice point. Thus, we must have $a_1 + \dots + a_k \geq 0$ for every $1 \leq k \leq 2n - 4$.

Theorem 7.2 (cf. Corollary 6.2.3 in [19]). *The Catalan number*

$$\text{Cat}_n = \frac{1}{n+1} \binom{2n}{n}, \quad (n \geq 0)$$

counts the number of sequences $a_1, \dots, a_{2n}$ of 1 's and -1 's with

$$a_1 + \dots + a_k \geq 0, \quad (k = 1, 2, \dots, 2n)$$

and $a_1 + \dots + a_{2n} = 0$.

Corollary 7.3. *The multiplicity of $X_{2,n}$ at P_σ is equal to the Catalan number*


$$\text{Cat}_{n-2} = \frac{1}{n-1} \binom{2n-4}{n-2}.$$

7.4. $\text{Mult}_{P_\tau} X_{2,n}$. In this section, we determine $\text{mult}_{P_\tau} X_{2,n}$, for τ being of block type (see Definition 7.7 below). Let τ be a face of σ , such that the associated (embedded sublattice) D_τ is of the form:

$$D_\tau = [(1, 2), (i, i + 1)] \cup [(i + k + 2, i + k + 3), (n - 1, n)] = I_1 \cup I_2, \quad \text{say,}$$

where $I_1 = [(1, 2), (i, i + 1)]$, $I_2 = [(i + k + 2, i + k + 3), (n - 1, n)]$, $1 \leq i \leq n - 3$, $0 \leq k \leq n - i - 3$.

We shall now determine $W(\tau)$ (cf. §6.7). Let A_τ denote the interval $[(1, i+2), (i+k+2, i+k+3)]$ in $J_{2,n}$:



Lemma 7.5. *With τ as above, we have $W(\tau) = \{e_{y'} - e_y \mid (y, y') \text{ is a cover in } A_\tau\}$.*

Proof. Clearly $e_{(n-1,n)}$ (the element in $W(\sigma)$ corresponding to the unique maximal element $(n-1, n)$ in $J_{2,n}$) is not in $W(\tau)$ (since $(n-1, n) \in D_\tau$). Let us denote

$$\theta = (i+k+2, i+k+3), \delta = (i, i+1)$$

Claim 1: For a cover (y, y') in A_τ , $f_{I_\alpha}(e_{y'} - e_y) = 0, \forall \alpha \in D_\tau$.

The claim follows in view of the facts that for a cover (y, y') in A_τ , we have,

- (i) $y, y' \in I_\theta$, and hence $y, y' \in I_\alpha, \forall \alpha \in I_2$.
- (ii) $y, y' \notin I_\delta$, and hence $y, y' \notin I_\alpha, \forall \alpha \in I_1$.

Claim 2: For a cover (y, y') in $J_{2,n}$ not contained in A_τ , there exists an $\alpha \in D_\tau$ such that $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

Note that a cover in $J_{2,n}$ is one of the following three types:

Type I: $((1, j), (1, j-1)), 3 \leq j \leq n$

Type II: $((j-1, j), (j-2, j-1)), 4 \leq j \leq n$

Type III: $((j-1, j), (1, j)), 3 \leq j \leq n$

Let now (y, y') be a cover not contained in A_τ .

If (y, y') is of Type I, then $(y, y') = ((1, j), (1, j-1))$, where either $j \leq i+2$ or $j \geq i+k+4$. Letting

$$\alpha = \begin{cases} (1, j-1), & \text{if } j \leq i+2 \\ (j-2, j-1), & \text{if } j \geq i+k+4 \end{cases}$$

we have, $\alpha \in D_\tau$ and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

If (y, y') is of Type II, then $(y, y') = ((j-1, j), (j-2, j-1))$, where either $j \leq i+2$ or $j \geq i+k+4$. Letting $\alpha = (j-2, j-1)$, we have, $\alpha \in D_\tau$ and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

If (y, y') is of Type III, then $(y, y') = ((j-1, j), (1, j))$, where either $j \leq i+1$ or $j \geq i+k+4$. Letting

$$\alpha = \begin{cases} (1, j), & \text{if } j \leq i+1 \\ (j-2, j), & \text{if } j \geq i+k+4 \end{cases}$$

we have, $\alpha \in D_\tau$ and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

The required result follows from claims 1 and 2. $\square$

Corollary 7.6. *With τ as in Lemma 7.5, we have*

$$\tau = \sigma_{i,1} \cup \sigma_{i+1,1} \cup \dots \cup \sigma_{i+k,1}.$$

Definition 7.7. We define a face τ as in Lemma 7.5 as a *J-block* (namely, τ is an union of consecutive $\sigma_{i,1}$'s).

Remark 7.8. Note that in general an union of faces need not be a face.

7.9. The Hibi variety $Z_{2,r}$. For an integer $r \geq 3$, let $\widetilde{I_{2,r}}$ denote the distributive lattice $I_{2,r} \setminus \{(1,2), (r-1,r)\}$. We define $Z_{2,r}$ to be the Hibi variety associated to $\widetilde{I_{2,r}}$. Note (cf. Proposition 4.3) that the cone associated to $Z_{2,r}$ has a set of generators consisting of $\{e_{y'} - e_y\}$, (y, y') being a cover in the sublattice $[(1,3), (r-1,r)]$ of $J_{2,r}$ (the set of join irreducibles of $I_{2,r}$). In view of Theorem 3.17, (2), we have

$$\text{mult}_{\mathbf{0}} Z_{2,r} = \text{mult}_{\mathbf{0}} X_{2,r} (= \text{Cat}_{r-2})$$

(Here, $\mathbf{0}$ denotes the origin.)

Theorem 7.10. *Let τ be a face of σ which is a “J-block” of $k+1$ consecutive $\sigma_{i,1}$'s (as in Definition 7.7). We have an identification of X_τ (the open affine piece of X_σ corresponding to τ) with $Z_{2,k+4} \times (K^*)^m$ where $m = \text{codim}_\sigma \tau (= 2(n-k) - 6)$*

Proof. In view of 1.6 and Proposition 4.6, we have

$\text{codim}_\sigma \tau = \dim X(D_\tau) = \#\{\text{elements in a maximal chain in } D_\tau\}$; from this it is clear that $\text{codim}_\sigma \tau$ equals $2(n-k) - 6$. Next, in view of Lemma 7.5 and §7.9, we obtain an identification of X_τ with $Z_{2,k+4} \times (K^*)^m$ (m being as in the Theorem). $\square$

Theorem 7.11. *Let τ be as in Theorem 7.10.*

- (1) *We have an identification of $TC_{P_\tau} X_\sigma$ with $Z_{2,k+4} \times (K)^m$ where $m = \text{codim}_\sigma \tau (= 2(n-k) - 6)$; in particular, $TC_{P_\tau} X_\sigma$ is a toric variety.*
- (2) *$\text{mult}_{P_\tau} X_{2,n} = \text{Cat}_{k+2} (= \frac{1}{k+3} \binom{2k+4}{k+2})$.*

Proof. X_τ being open in X_σ , we may identify $TC_{P_\tau} X_\sigma$ with $TC_{P_\tau} X_\tau$; assertion (1) follows from this in view of Theorem 7.10 (and §3.12).

Assertion (2) follows from (1) and Corollary 7.3. $\square$

7.12. A Product Formula. In this subsection, we give a product formula for $\text{mult}_{P_\tau} X_{2,n}$ where τ is a union of pairwise non-intersecting and non-consecutive *J*-blocks (see Remark 7.15 below).

Let τ be a face of σ , such that the associated (embedded sublattice) D_τ is of the form:

$$\begin{aligned} D_\tau = & [(1,2), (i_1, i_1+1)] \cup [(i_1+k_1+2, i_1+k_1+3), (i_2, i_2+1)] \\ & \cup [(i_2+k_2+2, i_2+k_2+3), (n-1, n)] = J_1 \cup J_2 \cup J_3, \text{ say} \end{aligned}$$

where $i_1 + k_1 + 1 < i_2$, and $J_1 = [(1, 2), (i_1, i_1 + 1)]$, $J_2 = [(i_1 + k_1 + 2, i_1 + k_1 + 3), (i_2, i_2 + 1)]$, $J_3 = [(i_2 + k_2 + 2, i_2 + k_2 + 3), (n - 1, n)]$.

Consider the following sublattices in $J_{2,n}$ (the set of join-irreducibles in $I_{2,n}$):

$$A = [(1, i_1 + 2), (i_1 + k_1 + 2, i_1 + k_1 + 3)], B = [(1, i_2 + 2), (i_2 + k_2 + 2, i_2 + k_2 + 3)].$$

Lemma 7.13. *With τ as above, we have $W(\tau) = \{e_{y'} - e_y \mid (y, y') \text{ is a cover in } A \cup B\}$.*

Proof. We proceed as in the proof of Lemma 7.5. As in that proof, we have $e_{(n-1,n)}$ is not in $W(\tau)$ (since $(n - 1, n) \in D_\tau$). Let us denote

$$\theta_1 = (i_1 + k_1 + 2, i_1 + k_1 + 3), \theta_2 = (i_2 + k_2 + 2, i_2 + k_2 + 3), \delta_1 = (i_1, i_1 + 1), \delta_2 = (i_2, i_2 + 1).$$

For any cover (y, y') in $A \cup B$, we clearly have $y, y' \in I_{\theta_2}$ and hence $y, y' \in I_\alpha, \forall \alpha \in J_3$; also, $y, y' \notin I_{\delta_1}$ and hence $y, y' \notin I_\alpha, \forall \alpha \in J_1$. Thus we obtain that

$$(*) \quad f_{I_\alpha}(e_{y'} - e_y) = 0, \forall \alpha \in J_1 \cup J_3$$

Next, if (y, y') is a cover in A , then $y, y' \in I_{\theta_1}$ and hence $y, y' \in I_\alpha, \forall \alpha \in J_2$; also, if (y, y') is a cover in B , then $y, y' \notin I_{\delta_2}$, and hence $y, y' \notin I_\alpha, \forall \alpha \in J_2$ (note that θ_1 (resp. δ_2) is the smallest (resp. largest) element in J_2). Thus we obtain that

$$(**) \quad f_{I_\alpha}(e_{y'} - e_y) = 0, \forall \alpha \in J_2$$

Now $(*)$ and $(**)$ imply the inclusion “ $\supseteq$ ”. We shall prove the inclusion “ $\subseteq$ ” by showing that if a cover (y, y') is not contained in $A \cup B$, then there exists an $\alpha \in D_\tau$ such that $f_{I_\alpha}(e_{y'} - e_y) \neq 0$. This proof again runs on similar lines as that of Lemma 7.5. Let then (y, y') be a cover in $J_{2,n}$ not contained in $A \cup B$. It would be convenient to introduce the following sublattices in $J_{2,n}$:

$$\begin{aligned} P &= [(1, 2), (i_1 + 1, i_1 + 2)] \\ Q &= [(1, i_1 + k_1 + 3), (i_2 + 1, i_2 + 2)] \\ R &= [(1, i_2 + k_2 + 3), (n - 1, n)]. \end{aligned}$$

We distinguish the following cases:

Case I: Let (y, y') be of type I (cf. proof of Lemma 7.5), say, $((1, j), (1, j - 1))$.

(i) If (y, y') is contained in P , then $j \leq i_1 + 2$. We let $\alpha = (1, j - 1)$. Note that $\alpha \in J_1$ and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

(ii) If (y, y') is contained in Q (resp. R), then $i_1 + k_1 + 4 \leq j \leq i_2 + 2$ (resp. $i_2 + k_2 + 4 \leq j \leq n$). We let $\alpha = (j - 2, j - 1)$. Note that $\alpha \in J_2$ (resp. J_3) and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

Case II: Let (y, y') be of type II, say, $((j - 1, j), (j - 2, j - 1))$.

Then, $3 \leq j \leq i_1 + 2$, $i_1 + k_1 + 4 \leq j \leq i_2 + 2$ or $i_2 + k_2 + 4 \leq j \leq n$ accordingly as (y, y') is contained in P , Q or R . We let $\alpha = (j - 2, j - 1)$. Note that $\alpha \in J_1, J_2$ or J_3 , accordingly as (y, y') is contained in P, Q or R ; and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

Case III: Let (y, y') be of type III, say, $((j - 1, j), (1, j))$.

(i) If (y, y') is contained in P , then $j \leq i_1 + 1$. We let $\alpha = (1, j)$. Note that $\alpha \in J_1$ and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$.

(ii) If (y, y') is contained in Q (resp. R), then $i_1 + k_1 + 4 \leq j \leq i_2 + 1$ (resp. $i_2 + k_2 + 4 \leq j \leq n$). We let $\alpha = (j - 2, j)$. Note that $\alpha \in J_2$ or J_3 , accordingly as (y, y') is contained in Q or R ; and $f_{I_\alpha}(e_{y'} - e_y) \neq 0$. $\square$

As an immediate consequence of Lemma 7.13 and Corollary 7.6, we have

Corollary 7.14. *Let τ be as in Lemma 7.13. Then $\tau = \tau_1 \cup \tau_2$, where*

$$\begin{aligned}\tau_1 &= \sigma_{i_1,1} \cup \dots \cup \sigma_{i_1+k_1,1}, \\ \tau_2 &= \sigma_{i_2,1} \cup \dots \cup \sigma_{i_2+k_2,1};\end{aligned}$$

Remark 7.15. We refer to a pair (τ_1, τ_2) of faces as in Corollary 7.14 as *non-intersecting J -blocks*.

Theorem 7.16. *Let $\tau = \tau_1 \cup \tau_2$, where τ_1 and τ_2 are two non-intersecting (and non-consecutive) J -blocks (cf. Corollary 7.14). We have an identification of X_τ (the open affine piece of X_σ corresponding to τ) with $Z_{2,k_1+4} \times Z_{2,k_2+4} \times (K^*)^m$ where $m = \text{codim}_\sigma \tau (= 2(n - k_1 - k_2) - 9)$*

Proof is similar to that of Theorem 7.11 (using Lemma 7.13).
As an immediate consequence, we have

Theorem 7.17. *Let $\tau = \tau_1 \cup \tau_2$, where τ_1 and τ_2 are two non-intersecting (and non-consecutive) J -blocks (cf. Corollary 7.14) Then*

- (1) *We have an identification of $TC_{P_\tau} X_\sigma$ with $Z_{2,k_1+4} \times Z_{2,k_2+4} \times (K^*)^m$ where $m = \text{codim}_\sigma \tau (= 2(n - k_1 - k_2) - 9)$; in particular, $TC_{P_\tau} X_\sigma$ is a toric variety.*
- (2) *$\text{mult}_{P_\tau} X_{2,n} = (\text{mult}_{P_{\tau_1}} X_{2,n}) \cdot (\text{mult}_{P_{\tau_2}} X_{2,n})$.*

Proof is similar to that of Theorem 7.11 (using Theorem 7.16).

Remark 7.18. It is clear that we can extend this multiplicative property to $\tau = \tau_1 \cup \dots \cup \tau_s$, a union of s pairwise non-intersecting, non-consecutive J -blocks.

8. A MULTIPLICITY FORMULA FOR $X_{d,n}$

In this section, we give a formula for $\text{mult}_{P_\sigma} X_{d,n}$. By Theorem 3.17, (2), we have that $\text{mult}_{P_\sigma} X_{d,n}$ equals the number of maximal chains in $I_{d,n}$. We give below an explicit formula for the number of maximal chains in $I_{d,n}$. Notice that the number of chains in $I_{d,n}$ from $(1, 2, \dots, d)$ to $(n - d + 1, \dots, n)$ is the same as the number of chains from $(0, 0, \dots, 0)$ to $(n - d, n - d, \dots, n - d)$ such that for any $(i_1, \dots, i_d)$ in the chain, we have $i_1 \geq i_2 \geq \dots \geq i_d \geq 0$. Now, set

$$\mu = (\mu_1, \mu_2, \dots, \mu_d) = (n - d, n - d, \dots, n - d). \quad (*)$$

For any $\lambda \vdash m$, let $f^\lambda = K_{\lambda, 1^m}$, i.e., the number of standard Young tableau of shape λ (cf. [19]).

Proposition 8.1 (cf. Proposition 7.10.3, [19]). *Let λ be a partition of m . Then the number f^λ counts the lattice paths $0 = v_0, v_1, \dots, v_m$ in $\mathbb{R}^l$ (where $l = l(\lambda)$) from the origin v_0 to $v_m = (\lambda_1, \lambda_2, \dots, \lambda_l)$, with each step a coordinate vector; and staying within the region (or cone) $x_1 \geq x_2 \geq \dots \geq x_l \geq 0$.*

Thus, for μ as described in $(*)$ above, we have that the number of maximal chains in $I_{d,n}$ is equal to f^μ .

Corollary 7.21.5 of [19] gives an explicit description of f^λ .

Proposition 8.2. *Let $\lambda \vdash m$. Then*

$$f^\lambda = \frac{m!}{\prod_{u \in \lambda} h(u)}.$$

The statement above refers to $u \in \lambda$ as a box in the Young tableau of λ , and $h(u)$ being the “hook length” of u . The hook length is easily defined as the number of boxes to the right, and below, of u , including u once.

Let us take, for example, $I_{3,6}$. Then $\mu = (3, 3, 3)$, and the Young tableau of shape μ with hook lengths given in their corresponding boxes is

5	4	3
4	3	2
3	2	1

Therefore

$$f^\mu = \frac{9!}{5 \cdot 4^2 \cdot 3^3 \cdot 2^2 \cdot 1} = 42.$$

In fact, in the scenario of $I_{d,n}$ our derived partition μ (given by $(*)$) will always be a rectangle; and we can deduce a formula for f^μ which does not require the Young tableau. The top left box of μ will always have hook length $(n - d) + d - 1 = n - 1$. Then, the box directly below it, and the box directly to the right of it will have length $n - 2$. For any box of μ , the box below and the box to the right will have hook length one less than that of the box with which we started.

Since the posets $I_{d,n}$, and $I_{n-d,n}$ are isomorphic, we can assume that $d \leq n - d$. Then we have $\prod_{u \in \mu} h(u) =$

$$(n - 1)(n - 2)^2 \cdots (n - d)^d (n - d - 1)^d \cdots (d)^d (d - 1)^{d-1} \cdots (2)^2 (1).$$

Thus we arrive at the following.

Theorem 8.3. *The multiplicity of $X_{d,n}$ at P_σ is equal to*

$$\frac{(d(n - d))!}{(n - 1)(n - 2)^2 \cdots (n - d)^d (n - d - 1)^d \cdots (d)^d (d - 1)^{d-1} \cdots (2)^2 (1)}.$$

9. CONJECTURES

In this section, we give two conjectures on the multiplicity at a singular point and one conjecture on $\text{Sing } X(\mathcal{L})$, $\mathcal{L}$ being the Bruhat poset of Schubert varieties in a minuscule G/P .

Guided by the phenomenon for $X_{2,n}$ as given by Theorem 7.17, we make the following conjecture.

Conjecture 1. There exists a multiplicative formula in $X_{d,n}$ similar to Theorem 7.17 for a face which has the form as an union of several disjoint (and non-consecutive) faces.

The generating set $W(\tau)$ of a face τ consists of $\{e_{y'} - e_y\}$, for certain covers (y, y') in $J(\mathcal{L})$ (assuming that $\hat{1} \in D_\tau$ so that $e_{\hat{1}}$ is not in $W(\tau)$). Thus $W(\tau)$ determines a sub set $H(\tau) := \bigcup H(\tau)_i$ of $J(\mathcal{L})$, such that $W(\tau)$ consists of all the covers in the $H(\tau)_i$'s. Thus, $H(\tau)$ for $\tau = \sigma_{ij}$ would be the diamond given in Remark 6.10. In §7.12, if $\tau = \tau_1 \cup \tau_2$ for τ_1, τ_2 a pair of non-consecutive, non-intersecting J -blocks, $H(\tau) = H(\tau_1) \dot{\cup} H(\tau_2)$.

Theorem 7.11 implies that $\text{mult}_{P_{\tau_1}} X_{2,n} = \text{mult}_{P_{\tau_2}} X_{2,n}$ if both τ_1 and τ_2 are J -blocks of the same length; in particular, $H(\tau_1)$ and $H(\tau_2)$ are isomorphic. Guided by this phenomenon we make the following conjecture.

Conjecture 2. For a face τ of any Hibi toric variety $X(\mathcal{L})$, $\text{mult}_{P_\tau} X(\mathcal{L})$ is determined by the poset $H(\tau)$. By this we mean that if τ, τ' are such that $H(\tau), H(\tau')$ are isomorphic posets, then the multiplicities of $X(\mathcal{L})$ at the points $P_\tau, P_{\tau'}$ are equal.


Remark 9.1. Toward generalizing Theorem 6.19 to other Hibi varieties, we will first explain how the lattice points μ_{ij}, λ_{ij} were chosen. Let θ, δ be two incomparable, join and meet irreducibles in $I_{d,n}$; say $\theta = (1, \dots, j, n+j+1-d, \dots, n)$ and $\delta = (i+1, \dots, i+d)$. Then $\theta \wedge \delta = \mu_{ij}$ and $\theta \vee \delta = \lambda_{ij}$. In view of Theorems 6.15 and 6.17, we have

In $X_{d,n}$, P_τ is a smooth point if and only if for every pair (θ, δ) of join and meet irreducibles, there is some $\alpha \in [\theta \wedge \delta, \theta \vee \delta]$ such that $P_\tau(\alpha)$, the α -th co-ordinate of P_τ is non-zero.

This is in fact the content of the conjecture of [11] (see [11], §11).

This suggests for us to look at such pairs of join-meet irreducibles in other distributive lattices and expect the components of the singular locus of the associated Hibi toric variety to be given by Theorem 6.19,(1) for the case of $I_{d,n}$. This is not true in general, however, as the following counter example shows.

9.2. Counter Example. Let $\mathcal{L}$ be the interval $[(1, 3, 4), (2, 5, 6)]$, a sublattice of $I_{3,6}$.



Notice that $\mathcal{L}$ has only one pair of join meet irreducibles, $(2, 3, 4)$ and $(1, 5, 6)$; and thus the corresponding interval $[\theta \wedge \delta, \theta \vee \delta]$ is the entire lattice. Therefore, if our result (Theorem 6.19,(1)) on the singular locus of G-H toric varieties were to generalize to other Hibi toric varieties, we have that any proper face is non-singular, since any face τ must correspond to an embedded sublattice D_τ , and naturally this sublattice will intersect the interval, which is just $\mathcal{L}$.

But this is not true! For example, let τ be the face of σ such that $D_\tau = \{(1, 5, 6)\}$. Then, the following set

$$\tau = C \langle e_{145} - e_{156}, e_{136} - e_{156}, e_{135} - e_{145}, e_{135} - e_{136}, e_{134} - e_{135} \rangle.$$

is a set of generators for τ . Clearly τ is not generated by the subset of a basis, and thus τ is a singular face (see Lemma 6.14).

Nevertheless, Theorem 6.19 holds for minuscule posets as described below.

Let G be semisimple, and P a maximal parabolic subgroup with ω as the associated fundamental weight. Let W (resp. W_P) be the Weyl group of G (resp. P). Then the Schubert varieties in G/P are indexed by W/W_P . Let P be *minuscule*, i.e., the weights in the fundamental representation associated to ω form one orbit under the Weyl group. One knows that the Bruhat poset W/W_P of the Schubert varieties in G/P is a distributive lattice. See [14] for details.

Definition 9.3. We call $\mathcal{L} := W/W_P$, a *minuscule poset*, and $X(\mathcal{L})$, a *Bruhat-Hibi toric variety*, or a B-H toric variety (for short).

Remark 9.4. Any G-H toric variety $X_{d,n}$ is a B-H toric variety.

Now for $\mathcal{L}$ as in Definition 9.3, consider a pair (α, β) of incomparable join-meet irreducible elements. Let $I_{\alpha, \beta} = [(\alpha \wedge \beta), (\alpha \vee \beta)]$.

It has been shown very recently (cf.[6]) that a B-H toric variety $X(\mathcal{L})$ is smooth at P_τ (τ being a face of σ) if and only if for each incomparable pair (α, β) of join-meet irreducibles in $\mathcal{L}$, there exists at least one $\gamma \in I_{\alpha, \beta}$ such that $P_\tau(\gamma)$ is non-zero.

REFERENCES

- [1] M. Aigner, *Combinatorial Theory*, Springer-Verlag, New York, 1979.
- [2] Victor V. Batyrev, Ionut Ciocan-Fontanine, Bumsig Kim, Duco van Straten, *Mirror Symmetry and Toric Degenerations of Partial Flag Manifolds*, Acta Math. 184 (2000), no. 1, 1–39.
- [3] Victor V. Batyrev, Ionut Ciocan-Fontanine, Bumsig Kim, Duco van Straten, *Conifold Transitions and Mirror Symmetry for Calabi-Yau Complete Intersections in Grassmannians*, Nuclear Phys. B 514 (1998), no. 3, 640–666.
- [4] A. Borel, *Linear Algebraic Groups*, second edition, Springer-Verlag, New York, 1991.
- [5] N. Bourbaki, *Groups et Algèbres de Lie*, Chapitres 4, 5 et 6, Hermann, Paris, 1968.
- [6] J. Brown and V. Lakshmibai, *Singular Loci of Bruhat-Hibi Toric Varieties*, to appear in J. Algebra.
- [7] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Springer-Verlag, New York, 1995.
- [8] D. Eisenbud and B. Sturmfels, *Binomial ideals*, Duke Mathematical Journal **84** (1996), 1–45. (1994).
- [9] W. Fulton, *Introduction to Toric Varieties*, Annals of Math. Studies **131**, Princeton U. P., Princeton N. J., 1993.
- [10] N. Gonciulea and V. Lakshmibai, *Degenerations of flag and Schubert varieties to toric varieties*, Transformation Groups, vol 1, no:3 (1996), 215-248.
- [11] N. Gonciulea and V. Lakshmibai, *Schubert varieties, toric varieties and ladder determinantal varieties*, Ann. Inst. Fourier, t.47, 1997, 1013-1064.
- [12] R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, New York, 1977.
- [13] T. Hibi, *Distributive lattices, affine semigroup rings, and algebras with straightening laws*, Commutative Algebra and Combinatorics, Advanced Studies in Pure Math. **11** (1987) 93-109.
- [14] H. Hiller, *Geometry of Coxeter groups*, Pitman Adv. pub. prog., Research Notes in Math, vol 54, Boston, London, Melbourne, 1982.
- [15] G. Kempf et al, *Toroidal Embeddings*, Lecture notes in Mathematics, N0. 339, Springer-Verlag, 1973.
- [16] V. Lakshmibai and N. Gonciulea, *Flag Varieties*, Hermann, Éditeurs des Sciences et des Arts, 2001.
- [17] V. Lakshmibai and H. Mukherjee, *Singular Loci of Certain Toric Varieties*, submitted to J. Ramanujan Math. Soc.
- [18] R.A. Proctor, *Bruhat Lattices, Plane Partition Generating Functions, and Minuscule Representations*, European J. Combinatorics **5** (1984), 331-350.
- [19] R.P. Stanley, *Enumerative Combinatorics Volume 2*, Cambridge University Press, 1999.
- [20] B. Sturmfels, *Gröbner bases and convex polytopes*, American Mathematical Society, Univ. Lecture Series, No. 8, Providence, Rhode Island, 2002.
- [21] D. G. Wagner, *Singularities of toric varieties associated with finite distributive lattices*, Journal of Algebraic Combinatorics **5** (1996), 149–165.

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115

E-mail address: brown.justin1@neu.edu, lakshmibai@neu.edu